

nitriCLEAR[™] Aloe AF

POWDER FREE NITRILE EXAMINATION GLOVES

Standards Met: ASTM D6319, ASTM D6978*

Attributes:

- Not made with natural rubber latex
- Not made with accelerators
- Ambidextrous
- Short beaded cuff
- Finger textured
- Low dermatitis potential
- Aloe Vera coating
- Non-sterile
- Single use only

Dimensions:


Box dimensions: 23.0 cm x 12.5 cm x 8 cm

Box count: 250 gloves per box (XS-L), 200 gloves per box (XL)

Case pack: 10 boxes per case



SPECIFICATIONS	X-SMALL	SMALL	MEDIUM	LARGE	X-LARGE
Reorder Number	#51490	#51491	#51492	#51493	#51494
Width of Palm (mm)	75 ± 5	85 ± 5	95 ± 5	106 ± 5	115 ± 5
Total Length (mm)	240 min.	240 min.	240 min.	240 min.	240 min.
Thickness: Cuff (mm)	0.04 min.	0.04 min.	0.04 min.	0.04 min.	0.04 min.
Thickness: Palm (mm)	0.05 min.	0.05 min.	0.05 min.	0.05 min.	0.05 min.
Thickness: Finger (mm)	0.08 min.	0.08 min.	0.08 min.	0.08 min.	0.08 min.
Tensile Strength (MPa)	14 min.	14 min.	14 min.	14 min.	14 min.
Elongation (%) Before Aging	500 min.	500 min.	500 min.	500 min.	500 min.
Elongation (%) After Aging	400 min.	400 min.	400 min.	400 min.	400 min.



*Tested for use with Chemotherapy Drugs per **ASTM D6978: Standard Practice For Assessment Of Resistance Of Medical Gloves To Permeation By Chemotherapy Drugs**. Gloves used for protection against chemotherapy drug exposure should be selected specifically for the type of chemicals used. Users are recommended to review Safety Data Sheets for Chemotherapy Drugs being used to determine the required level of protection.



DO NOT USE WITH CARMUSTINE OR THIOTEPA. Testing showed an average breakthrough time of 11 minutes with Carmustine and 59 minutes with Thiotepea.

*Tested Chemotherapy Drug and Concentration	Average Breakthrough Detection Time (minutes)
Carboplatin, 10 mg/mL	No breakthrough up to 240 minutes
Cisplatin (BCNU), 1.0 mg/mL	No breakthrough up to 240 minutes
Cyclophosphamide (Cytoxan), 20 mg/mL	No breakthrough up to 240 minutes
Dacarbazine (DTIC), 10 mg/mL	No breakthrough up to 240 minutes
Doxorubicin Hydrochloride, 2.0 mg/mL	No breakthrough up to 240 minutes
Etoposide (Toposar), 20 mg/mL	No breakthrough up to 240 minutes
Fluorouracil, 50 mg/mL	No breakthrough up to 240 minutes
Ifosfamide, 50 mg/mL	No breakthrough up to 240 minutes
Methotrexate, 25 mg/mL	No breakthrough up to 240 minutes
Mitomycin C, 0.5 mg/mL	No breakthrough up to 240 minutes
Mitoxantrone, 2.0 mg/mL	No breakthrough up to 240 minutes
Paclitaxel (Taxol), 6.0 mg/mL	No breakthrough up to 240 minutes
Vincristine Sulfate, 1.0 mg/mL	No breakthrough up to 240 minutes
The following chemotherapy drugs and concentrations showed breakthrough detected in less than 240 minutes:	
Carmustine, 3.3 mg/mL	11 minutes
Thiotepea, 10 mg/mL	59 minutes

Made in Malaysia

The information provided is given in good faith and believed to be accurate, but no warranty or guarantee is given or implied.

MP-REF-51490B
Rev.2: 2022-05-11