

ACTIVE INGREDIENTS:

4.73 L

CAUTION KEEP OUT OF REACH OF CHILDREN READ THE LABEL BEFORE USING

PRECAUTIONARY STATEMENTS

Hazard to human and domestic animals. Do not breathe vapor. Causes moderate eye irritation. Avoid contact with eyes, or clothing. Wear protective eye wear and gloves. Wash thoroughly with soap and water after handling.

REV.2019/09

tb Minuteman NEX GEN Disinfectant is intended for use on pre-cleaned, hard non-norous inanimate surfaces in areas where the disinfection of surfaces is required. When used as directed to Minuteman NEX GEN Disinfectant cleans and disinfects hard, non-porous, non-food contact surfaces found in ambulances, dental facilities, hospitals, nursing homes, paramedic units, schools, clinics, parlors, bathrooms or any other places that are concerned with the hazards of cross-contamination from infectious microorganisms. When used as directed tb Minuteman NEX GEN Disinfectant may be used on the following hard, non-porous, non-food contact surfaces: bed & security rails, cabinets, commodes, counters, equipment surfaces & knobs, pet care surfaces, tables, toys, vehicle interior surfaces. When used as directed tb Minuteman NEX GEN Disinfectant may be used on the following hard, non-norous, non-critical parts of the following items/surfaces; ambulance equipment, blood pressure devices, CPR training devices, medical equipment and dental chairs# EMS equipment, exam tables, respiratory care & therapy equipment and any surfaces that are potentially contaminated with disease-causing microorganisms. If you are uncertain as to the material composition of your surface, confirm compatibility with the manufacturer before proceeding. For plastic and painted surfaces, spot test on an inconspicuous area before use.

DIRECTIONS FOR USE

This product is not to be used as a terminal sterilant high-level disinfectant on any surface or instrument that (i) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean critical or semi-critical medical devices prior to sterilization/high-level disinfection.

Placing the Contents of the Refill Jug into an Empty Sprayer:

Remove the cap and bottle seal. Discard seal and retain the cap.
 Pour product from the refill jug into the empty and clean sprayer bottle, being careful not to overfill the sprayer bottle.



3. Insert and secure the sprayer to its bottle.4. Secure the retained cap snugly to the refill jug.

CLEANING AND DISINFECTING

Pre-clean any heavily soiled surfaces. Spray the surface with the Minuteman NEX GEN Disinfectant to pre-clean the surface. Using a disposable cloth, clean the surface. Discard. Spray the Minuteman NEX GEN Disinfectant about 20 cm from the surface, thoroughly wetting area to be disinfected. Treated surface must remain visibly wet for the appropriate time indicated for the purpose intended. For broad spectrum disinfection, allow the surface to remain wet for 3 minutes. Then, wipe the surface to remain wet for 3 minutes. Then, wipe the surface to remain.

Do not contaminate food during use and storage.

A potable water rinse is recommended after disinfection if:

- any surfaces may come into direct contact with children at the
mouthing stage, or

- any surfaces may come into contact with food or drinking water.

Made in Canada with domestic and imported ingredients

riade in Canada with domestic and imported ingredient

St. Thomas, ON, Canada, N5R 6C8 www.maxill.com 1-800-268-8633

 Questions?
 Call 1-800-268-8633, 8 am - 5 pm ET Weekdays

 For Emergencies, call CANUTEC at 613-996-6666
 SDS No. MP-RFF-61160S

REV 2019/09

ORGANISMS CONTROLLED / CONTACT TIMES

When used as directed on pre-cleaned hard, non-porous non-food contact surfaces, tb Minuteman NEX GEN Disinfectant disinfects the following microorganisms with a contact time:

1-minute contact time

Human Immunodeficiency virus type 1 (HIV-1)* Duck Hepatitis B Virus (Surrogate for Human Hepatitis B Virus)* Bovine Viral Diarrhea Virus

(Surrogate for Human Hepatitis C Virus)*

Rotavirus

Human Coronavirus Avian Influenza A (H3N2) Reassortant virus

Herpes Simplex Virus Type 1 Herpes Simplex Virus Type 2

3-minute contact time

Mycobacterium bovis BCG Pseudomonas aeruginosa Staphylococcus aureus Salmonella enterica

Vancomycin Resistant Enterococcus faecalis (VRE) Methicillin Resistant Staphylococcus aureus (MRSA) Adenovirus type 5 (ATCC VR-5)

Feline Calicivirus (Surrogate for Human Norovirus)
Trichophyton mentagrophytes

"This product effectively kills Human Immunodeficiency virus, ype I (IRIV-). Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV) on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects can be associated with the optential for transmission of HIV-1 HBV and HCV.

Special Decontamination Procedures:

Personal protection: Personnel that clean items soiled with blood or body fluids must wear appropriate personal protective equipment (i.e. barrier protection), such as disposable gloves, gowns, masks or eye coverings;

Cleaning: Target surfaces must be cleaned prior to disinfection, as the presence of heavy soil on a surface may reduce the intended disinfectant efficacy of the product; and Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1. HBV and HCV.

Infectious materials disposal: Any materials used in the cleaning process that may contain blood or body fluids are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

FIRST AID: If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give CPR. If in eyes, hold eye open and rinses slowly and gently with water for IS - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then confluer in Call a poison control center or doctor for treatment advice. Have the product table and/or Drug I dehrification Number (DIN) with you when calling a poison control center, doctor, or opin for treatment.

PHYSICAL HAZARDS: Combustible. Do not use or store near heat or open flame. STORAGE: Store in a cool place. Do not reuse this container.

DISPOSAL: Recycle this container or discard container in trash. For information on disposal of unused, unwanted product, contact the manufacturer or the appropriate municipal/provincial/territorial agency or product stewardship organization.



Cleaning Care for Dental Chairs

In addition to disinfecting, a cleaning protocol is required by the chair manufacturer. Call **1-800-268-8633** for more information. **www.maxill.com/ca/chaircare**