

tb Minuteman NEX GEN Disinfectant is intended for use on pre-cleaned, hard non-porous inanimate surfaces in areas where the disinfection of surfaces is required.

When used as directed tb Minuteman NEX GEN Disinfectant cleans and disinfects hard, non-porous, non-food contact surfaces found in ambulances, dental facilities, hospitals, nursing homes, paramedic units, schools, clinics, parlors, bathrooms or any other places that are concerned with the hazards of cross-contamination from infectious microorganisms.

When used as directed tb Minuteman NEX GEN Disinfectant may be used on the following hard, non-porous, non-food contact surfaces: bed & security rails, cabinets, commodes, counters, equipment surfaces & knobs, pet care surfaces, tables, toys, vehicle interior surfaces.

When used as directed tb Minuteman NEX GEN Disinfectant may be used on the following hard, non-porous, non-critical parts of the following items/surfaces: ambulance equipment, blood pressure devices, CPR training devices, medical equipment and dental chairs[†], EMS equipment, exam tables, respiratory care & therapy equipment and any surfaces that are potentially contaminated with disease-causing microorganisms. If you are uncertain as to the material composition of your surface, confirm compatibility with the manufacturer before proceeding. For plastic and painted surfaces, spot test on an inconspicuous area before use.

ORGANISMS CONTROLLED / CONTACT TIMES[†]

When used as directed on pre-cleaned hard, non-porous non-food contact surfaces, tb Minuteman NEX GEN Disinfectant disinfects the following microorganisms with a *contact time*:

1-minute contact time

Human Immunodeficiency virus type 1 (HIV-1)*
Duck Hepatitis B Virus (Surrogate for Human Hepatitis B Virus)*
Bovine Viral Diarrhea Virus (Surrogate for Human Hepatitis C Virus)*
Rotavirus
Human Coronavirus
Avian Influenza A (H3N2) Reassortant virus
Herpes Simplex Virus Type 1
Herpes Simplex Virus Type 2

3-minute contact time

Mycobacterium bovis BCG
Pseudomonas aeruginosa
Staphylococcus aureus
Salmonella enterica
Vancomycin Resistant *Enterococcus faecalis* (VRE)
Methicillin Resistant *Staphylococcus aureus* (MRSA)
Adenovirus type 5 (ATCC VR-5)
Feline Calicivirus (Surrogate for Human Norovirus)
Trichophyton mentagrophytes

FIRST AID: If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give CPR. If in eyes, hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. Have the product label and/or Drug Identification Number (DIN) with you when calling a poison control center, doctor, or going for treatment.

PHYSICAL HAZARDS: Combustible. Do not use or store near heat or open flame.

STORAGE: Store in a cool place. Do not reuse or refill except as described in the directions for use. Refill only with this product.

DISPOSAL: Recycle this container or discard container in trash. For information on disposal of unused, unwanted product, contact the manufacturer or the appropriate municipal/provincial/territorial agency or product stewardship organization.

maxill



Minuteman NEX GEN DISINFECTANT

MYCOBACTERICIDAL • VIRUCIDAL[†] • BACTERICIDAL • FUNGICIDAL
INTERMEDIATE LEVEL HOSPITAL DISINFECTANT SPRAY

Reorder / Item Number:
61155



ACTIVE INGREDIENTS:

n-Alkyl (50% C14, 40% C12, 10% C16)
dimethyl benzyl ammonium chloride 0.34% w/w
Didecyl dimethyl ammonium chloride 0.51% w/w
Ethanol 6.84% w/w
Isopropanol 13.68% w/w

DIN 02488981

946 mL

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ THE LABEL BEFORE USING

PRECAUTIONARY STATEMENTS

Hazard to human and domestic animals. Do not breathe vapor. Causes moderate eye irritation. Avoid contact with eyes, or clothing. Wear protective eye wear and gloves. Wash thoroughly with soap and water after handling.

DIRECTIONS FOR USE

This product is not to be used as a terminal sterilant/ high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean critical or semi-critical medical devices prior to sterilization/high-level disinfection.

Setting up the Sprayer:

1. Remove the cap and bottle seal. Discard.
2. Insert and secure the provided sprayer to the bottle.
3. Keep upright.

This container may be refilled by the consumer with tb Minuteman NEX GEN Disinfectant sold in jug quantities. Refer to further instructions on the label of the jug.

CLEANING AND DISINFECTING

Pre-clean any heavily soiled surfaces. Spray the surface with tb Minuteman NEX GEN Disinfectant to pre-clean the surface. Using a disposable cloth, clean the surface. Discard. Spray tb Minuteman NEX GEN Disinfectant about 20 cm from the surface, thoroughly wetting area to be disinfected. Treated surface must remain visibly wet for the appropriate time indicated for the purpose intended. For broad spectrum disinfection, allow the surface to remain wet for 3 minutes. Then, wipe the surface dry.

Do not contaminate food during use and storage.

A potable water rinse is recommended after disinfection if:

- any surfaces may come into direct contact with children at the mouthing stage, or
- any surfaces may come into contact with food or drinking water.

[†]This product effectively kills Human Immunodeficiency virus type 1 (HIV-1), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV) on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects can be associated with the potential for transmission of HIV-1, HBV, and HCV.

Special Decontamination Procedures:

Personal protection: Personnel that clean items soiled with blood or body fluids must wear appropriate personal protective equipment (i.e. barrier protection), such as disposable gloves, gowns, masks or eye coverings;

Cleaning: Target surfaces must be cleaned prior to disinfection, as the presence of heavy soil on a surface may reduce the intended disinfectant efficacy of the product; and

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1, HBV and HCV.

Infectious materials disposal: Any materials used in the cleaning process that may contain blood or body fluids are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

[†]Cleaning Care for Dental Chairs

In addition to disinfecting, a cleaning protocol is required by the chair manufacturer. Call 1-800-268-8633 for more information. www.maxill.com/ca/chaircare

Made in Canada with domestic and imported ingredients

maxill inc.
St. Thomas, ON, Canada, N5R 6C8
www.maxill.com 1-800-268-8633

Questions? Call 1-800-268-8633, 8 am - 5 pm ET Weekdays
For Emergencies, call CANUTEC at 613-996-6666
SDS No. MP-REF-611B03 REV.2019/09